

A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI

[\[Sign In\]](#) [\[Register\]](#)[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Books](#)Search  for   ☒ Limits☐ Preview/Index☐ History☐ Clipboard☐ DetailsDisplay  Show  Sort by  Send to 

All: 1

Review: 0

☐ 1: [Genes Dev.](#) 1993 Nov;7(11):2220-34.[Links](#)**A single cDNA, hTFIIA/alpha, encodes both the p35 and p19 subunits of human TFIIA.****DeJong J, Roeder RG.**

Laboratory of Biochemistry and Molecular Biology, Rockefeller University, New York, New York 10021.

TFIIA is a transcription factor that, by interacting with the TATA-binding subunit (TBP) of TFIID, modulates transcription initiation by RNA polymerase II in vitro. By use of a mobility shift assay, TFIIA was purified from HeLa cells as a complex of 35-, 19-, and 12-kD subunits. Oligonucleotides were used to isolate a human cDNA clone, hTFIIA/alpha, which encodes a 55-kD protein with homology to the product of the yeast gene TOA1. The open reading frame of hTFIIA/alpha contains peptide sequences obtained from both the p35 and p19 subunits of natural human TFIIA, and thus encodes these two subunits. Consistent with this, antiserum raised against the 55-kD hTFIIA/alpha-encoded protein reacted with both the p35 and p19 subunits of natural TFIIA, and the recombinant protein could functionally replace those subunits in a mobility shift assay with renatured p12. An efficient affinity purification for natural human TFIIA was suggested by the sequence of the hTFIIA/alpha protein and demonstrated biochemically. Finally, transcription from the adenovirus major late promoter was greatly reduced in nuclear extracts depleted with anti-TFIIA/alpha serum and was restored to original levels by the readdition of purified human TFIIA.

PMID: 8224848 [PubMed - indexed for MEDLINE]

Display  Show  Sort by  Send to [Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

### Related Links

Human general transcription factor TFIIA: characterization of a cDNA encoding the small subunit and requirement for basal and activated transcription. [Proc Natl Acad Sci U S A. 1995]

Molecular cloning of the small (gamma) subunit of human TFIIA reveals functions critical for activated transcription. [Genes Dev. 1994]

Characterization of the highly conserved TFIIA small subunit from Drosophila melanogaster. [Mol Cell Biol. 1994]

Isolation of a cDNA encoding the largest subunit of TFIIA reveals functions important for activated transcription. [Genes Dev. 1993]

Analysis of the yeast transcription factor TFIIA: distinct functional regions and a polymerase II-specific role in basal and activated transcription. [Mol Cell Biol. 1995]

[See all Related Articles...](#)

Feb 7 2007 15:04:00